Lake Overview


Monitoring began at Clark Lake in 2003. The data collected classify this lake in the city of Kent at moderate to high in primary productivity (mesotrophic - eutrophic) with fair water quality. Since the lake surface makes up only 2% of the 338-acre catchment basin, direct precipitation is less important than stormwater runoff and groundwater inputs. There is a 46 acre Class 1 wetland, which includes the lake and extends west into land historically used as pasture (King County, 1990).

Clark Lake is included in a Kent city park and is open to car top boats. City staff and lake users should watch nearshore aquatic plants to catch early infestations of Eurasian milfoil, Brazilian elodea or other noxious aquatic weeds.


Physical Parameters

The Level II Secchi transparency ranged between 1.4 and 2.2m during the sampling season. The May through October surface water temperatures reached a maximum of 26.6 degrees Celsius. No precipitation or water level records were available for the year.

Lake Temperature



Secchi Depth




Lake Level and Precipitation


No Data Available

Clark


Nutrient Analysis



TSI Ratings



Chlorophyll a Concentrations (ug/L)



Common algae Group

	•			
Anabaena sp.	bluegreen			
Cryptomonas sp.	cryptophyte			
Dinobryon sp.	chrysophyte			

Nutrient Analysis and TSI Ratings

Total phosphorus remained fairly constant through the season, with one anomalous high value in May, while total nitrogen decreased slowly from May through July and then rose again in September. The N:P ratio ranged from 13 to 52, suggesting occasionally good conditions for bluegreens. The 2003 TSI values for the three indicators straddled the line between mesotrophic and eutrophic, with TSI-Secchi higher than the other two indicators. This higher value may be affected by water color.

Chlorophyll and Algae

Chlorophyll content peaked in late June and then rose again in fall. The algae were dominated in the spring peak by *Cryptomonas* species and in fall by the bluegreen *Anabaena*. The chrysophyte *Dinobryon* was also important in the phytoplankton.

Clark 2003 Level I Data

No Level I Data Available For This Lake

Clark 2003 Level II Data

Date	Temp	Secchi	Chl-a	TP	TN	_	N:P	Calculated TSI			
(2003)	(°C)	(m)	(μg/l)	(μg/l)	(μg/l)	Obsv.		Secc	chl-a	TP	Notes
21-Apr											No Sample. Began
											monitoring 5-May-03.
5-May	14.0	2.1	3.5	18.7	980	1	52	49.3	42.8	46.4	
40 May	45.4	0.0	4.0	74.0	000	4	40	50.0	440	00.0	
19-May	15.4	2.0	4.3	74.0	936	1	13	50.0	44.9	66.2	
2-Jun	20.0	2.1	3.4	16.5	818	1	50	49.3	42.6	44.6	
2-3011	20.0	2.1	J. 4	10.5	010	'	30	49.5	42.0	44.0	
16-Jun	19.0	1.7	4.8	17.6	702	2	40	52.3	46.0	45.5	Chlor-a analysis error.
10 0411	10.0	'.,	1.0	17.0	.02	_		02.0	10.0	10.0	Estimated Chlor-a is 4.8 ug/L
30-Jun	22.0	1.5	16.0	19.4	690	2	36	54.1	57.8	46.9	-
14-Jul	25.5	1.5	6.4	19.9	619	1	31	54.1	48.8	47.3	
28-Jul	26.6	1.5	3.5	18.6	716		38	54.1	42.9	46.3	
11-Aug	22.6	2.2	2.7	16.1	572	1	36	48.6	40.2	44.2	
25-Aug	22.4	2.2	3.6	15.1	642	2	43	48.6	43.1	43.3	
0.0	04.5	4.5	40.0	05.0	070	0	00	F4.4	50.4	- 4 4	
8-Sep	21.5	1.5	16.6	25.9	672	3	26	54.1	58.1	51.1	
22-Sep											No sample.
22-3 6 p											ino sample.
7-Oct	16.9	1.4	14.9	16.1	811	3	50	55.1	57.1	44.2	
, 551	10.0				0			00.1	0		
20-Oct											No sample.
											·
	Temp	Secchi	Chl-a	TP	TN	Algae	N:P	Calculated TSI		TSI	
	(°C)	(m)	(μg/l)	(μg/l)	(μg/l)	Obsv.		Secc	chl-a	TP	
Mean	20.5	1.8	7.2	23.4	741.6	1.7	38	51.8	47.7	47.8	TSI Average = 49.2
Median	21.5	1.7	4.3	18.6	702.0	2	38	52.3	44.9	46.3	
Min	14.0	1.4	2.7	15.1	572.0	1	13	48.6	40.2	43.3	
Max	26.6	2.2	16.6	74.0	980.0	3	52	55.1	58.1	66.2	
Count	11	11	11	11	11	10	11	11	11	11	